

Siliguri Lab: Vive Health Care, C/o. Dr. P. D. Bhutia, Nirmal Nagar, Khaprail More, P.O. Matigara - 734010, Siliguri, Dt: Darjeeling Mobile: 7699991171, E-mail: siliguri.unipath@gmail.com

TEST REPORT

Reg. No: 18092700532 Reg. Date: 12-Sep-2018 00:00

Collected On: 12-Sep-2018 12:36

Name: **BISHNU SAIBA** Report Date: 12-Sep-2018

42 Years Sex: Male Dispatch At:

Age:

DR. P.D.BHUTIA MD.

Tele No:

9679791157

Ref. By: Location: Parameter

Result

Unit

Biological Reference Interval

HUMAN IMMUNODEFICIENCY VIRUS I & II

HIV I & II, Serum

Non Reactive

Flow-throgh Immunochromatography

Notes:

- 1) A NON REACTIVE result implies that no Anti HIV-1 or HIV-2 antibodies have been detected in the sample by this method. This means that either the patient has not been exposed to HIV-1 or HIV-2 infection or the sample has been tested during the "Window phase" (before the development of detectable levels of antibodies).
- 2) A REACTIVE result suggests the possibility of HIV-1 or /and HIV-2 infection. However these results must be verified by a confirmatory WESTERN BLOT method before declaring the patient positive for HIV-1 or HIV-2 infection.
- 3) Very high levels of IgM antibodies or anti-HLA A,B,C,DR antibodies can give false positive reaction.
- 4) Neonates born of HIV infected mothers may have HIV infection or can be uninfected despite the presence of maternal antibodies to HIV in their blood. Such neonates should undergo additional testing such as Polymerase chain reaction (PCR) to ascertain their status of
- 5) Pre & post test counseling for HIV testing is responsibility of referring physician.

HEPATITIS B SURFACE ANTIGEN

HBsAg, Serum

Negative

Negative

Immunochromatography

ANTI HCV

ANTI HCV, Serum

Non Reactive

Flow-through Immunochromatography

This is an electronically authenticated report.

S. Brance

* Denotes Test not in NABL Scope.

Approved by:

DR. SANKARSAN BHADURI MBBS, MD

(PATHOLOGY)

Approved On:

12-Sep-2018 16:50

Generated On: 12-Sep-2018 16:56